



Characterization of j -open sets in minimal structure spaces

M. Deepa^{1*} and S.S. Surekha²

Abstract

The aim of this paper is to introduce the notion of m - j open sets in minimal structure spaces together with its corresponding interior and closure operators. Furthermore, we investigated the basic properties of the sets and studied their relationship with other existing sets. Moreover, the minimal structure properties of j -border, j -frontier and j -exterior of the sets have been discussed.

Keywords

m - j open; m - j closed; m - j border; m - j frontier and m - j exterior.

AMS Subject Classification

54BXX, 54FXX.

^{1,2}Department of Mathematics, PSGR Krishnammal College for Women, Coimbatore-641004, Tamil Nadu, India.

*Corresponding author: ¹ mdeepa@psgrkcw.ac.in ²surekhasuresh97@gmail.com

Article History: Received 01 June 2020; Accepted 04 July 2020

©2020 MJM.

Contents

1	Introduction	1171
2	Preliminaries	1171
3	m - j open sets	1172
4	m - j border, m - j frontier, m - j exterior of the sets... ..	1173
	References	1174

1. Introduction

Popa and Noiri [12] introduced the concepts of minimal structure (briefly m -structure) in 2000. Also they studied the notion of m_x -open and m_x -closed sets and distinguish the sets using m_x -closure and m_x -interior operators respectively. The space which contains the mother set X and the null set \emptyset is said to be minimal structure. In 2009, W.K.Min and Y.K. Kim introduced Semiopen[7], Preopen[8] and α -open[9] in minimal structure. In 2012, I.Arockiarani and D.Sasikala[2] introduced the notion of j -open sets. In [6], Caldas studied the properties of α -border, α -frontier and α -exterior of α -open sets in topological space. In this paper, we introduced and studied the minimal structure properties of m - j border, m - j frontier and m - j exterior of the j -open sets. Also, we discussed some of the basic properties of m - j open sets in minimal structure.

2. Preliminaries

Definition 2.1. [12] Let X be a non-empty set and let $m_x \subseteq P(X)$ denotes the power set of X . We say that m_x is an minimal structure on X if \emptyset and X belongs to the minimal structure m_x . The sets which are in minimal structure are called m -open sets and the complement of the m -open sets are called m -closed sets.

Definition 2.2. [12] Let A be the subset of X . The m_x -interior and m_x -closure of the subset A is defined by
 i) m_x -Int(A) = $\cup\{U : U \in m_x; U \subseteq A\}$; where U is an open set.
 ii) m_x -Cl(A) = $\cap\{F - A \subseteq F; X - F \in m_x\}$, where F is a closed set.

Definition 2.3. Let X be the non-empty set and A be the subset of X . Then A is called

- i) m -semi open [7] if $A \subseteq m_x$ -Cl(m_x -Int(A))
- ii) m -preopen [8] if $A \subseteq m_x$ -Int(m_x -Cl(A))
- iii) m - b open [10] if $A \subseteq m_x$ -Int(m_x -Cl(A)) \cup m_x -Cl(m_x -Int(A))
- iv) m - α open [9] if $A \subseteq m_x$ -Int(m_x -Cl(m_x -Int(A)))

Lemma 2.4. Consider a minimal structure (X, m_x) . For the subsets A and B the following conditions are true:

- i) m_x -Cl($X - A$) = $X - m_x$ -Int(A) and m_x -Int($X - A$) = m_x -Cl(A)
- ii) If $(X - A) \in m_x$, then m_x -Cl(A) = A and if $A \in m_x$, then m_x -Int(A) = A
- iii) m_x -Cl(\emptyset) = \emptyset , m_x -Cl(X) = X , m_x -Int(\emptyset) = \emptyset , m_x -Int(X) = X

- iv) If $A \subset B$, then $m_x\text{-Cl}(A) \subset m_x\text{-Cl}(B)$ and $m_x\text{-Int}(A) \subset m_x\text{-Int}(B)$
 v) $A \subset m_x - \text{Cl}(A)$ and $m_x - \text{Int}(A) \subset A$
 vi) $m_x\text{-Cl}(m_x\text{-Cl}(A)) = m_x\text{-Cl}(A)$

Definition 2.5. [12] Let (X, m_x) be the minimal structure and A be the subset of X . A point $x \in X$ is called a limit point of A if and only if every neighborhood of A contains a point of A other than x and it is denoted by $D(A)$.

3. m - j open sets

Definition 3.1. Let (X, m_x) be the minimal structure. Let A be the subset of X . The subset A is called the m - j open set if $A \subseteq m_x\text{-Int}(m_x\text{-pcl}(A))$.

Example 3.2. Consider $X = \{p, q, r\}$ with minimal structure $m_x = \{\emptyset, X, \{p\}, \{r\}\}$. Then its corresponding m - j open sets are $\emptyset, X, \{p\}, \{q\}, \{r\}, \{p, r\}$.

Definition 3.3. Let (X, m_x) be the minimal structure with subset A of X . Then,

- i) The $m_x\text{-Int}_j(A)$ is the union of all the m - j open sets contained in A .
 ii) The $m_x\text{-Cl}_j(A)$ is the intersection of all the m - j closed sets containing A .

Theorem 3.4. Every m -open sets are m - j open sets.

Proof. Let (X, m_x) be the minimal structure. Let P be an m -open set, then we have $P = m_x\text{-Int}(P)$. Also we know that $P \subseteq m_x\text{-pcl}(P)$. Hence we get $P \subseteq m_x\text{-Int}(m_x\text{-pcl}(P))$. Therefore, P is m - j open set.

The converse of the above theorem is not true by the following example: Consider $X = \{p, q, r\}$ with minimal structure $m_x = \{\emptyset, X, \{p\}, \{q\}, \{r\}, \{p, r\}\}$. The set $A = \{p, r\}$ is m - j open but not m -open set. \square

Theorem 3.5. For any subsets A and B with minimal structure the following conditions are true. i) The $m_x\text{-Int}_j(A)$ is the largest m - j open set contained in A .

- ii) A is an m - j open set if and only if $A = m_x\text{-Int}_j(A)$.
 iii) $A \subseteq B$, then $m_x\text{-Int}_j(A) \subseteq m_x\text{-Int}_j(B)$.
 iv) $m_x\text{-Int}_j(\emptyset) = \emptyset$.
 v) $m_x\text{-Int}_j(X) = X$.
 vi) $m_x\text{-Int}_j(A \cap B) \subseteq m_x\text{-Int}_j(A) \cap m_x\text{-Int}_j(B)$.
 vii) $m_x\text{-Int}_j(A) \cup m_x\text{-Int}_j(B) \subseteq m_x\text{-Int}_j(A \cup B)$.

Proof. i) Since, $m_x\text{-Int}_j(A) = \cup\{G : G \text{ is } m\text{-}j \text{ open, } G \subset A\}$. Hence it contains every m - j open subset G of A . Therefore, $m_x\text{-Int}_j(A)$ is the largest m - j open set contained in A .

ii) Let $A = m_x\text{-Int}_j(A)$. Since $m_x\text{-Int}_j(A)$ is an m - j open set, A is also an m - j open set. Conversely, let A be any m - j open set and also it is the largest m - j open set. Hence we have $A = m_x\text{-Int}_j(A)$.

iii) Let $A \subseteq B$. We have to prove that $m_x\text{-Int}_j(A) \subseteq m_x\text{-Int}_j(B)$. Let $x \in m_x\text{-Int}_j(A)$. Then there exists an open set G such that $x \in G \subset A$. Since $A \subseteq B$, G is also contained in B . Hence

$x \in G \subset B$ which implies that $x \in m_x\text{-Int}_j(B)$. Hence $m_x\text{-Int}_j(A) \subseteq m_x\text{-Int}_j(B)$.

vi) Let $x \in m_x\text{-Int}_j(A \cap B)$. Hence there exist a open set G in X such that $x \in A \cap B$ which implies that $x \in A$ and $x \in B$. Therefore, $x \in G \subset A$ and $x \in G \subset B$. Hence $x \in m_x\text{-Int}_j(A)$ and $x \in m_x\text{-Int}_j(B)$. Hence $m_x\text{-Int}_j(A \cap B) \subseteq m_x\text{-Int}_j(A) \cap m_x\text{-Int}_j(B)$.

vii) Let $x \in m_x\text{-Int}_j(A) \cup m_x\text{-Int}_j(B)$. Hence $x \in m_x\text{-Int}_j(A)$ or $x \in m_x\text{-Int}_j(B)$. If $x \in m_x\text{-Int}_j(A)$, then there exists an open set G such that $x \in G \subset A$. If $x \in m_x\text{-Int}_j(B)$, then there exists an open set G such that $x \in G \subset B$. Hence $x \in G \subset A \cup B$. Therefore, $m_x\text{-Int}_j(A) \cup m_x\text{-Int}_j(B) \subseteq m_x\text{-Int}_j(A \cup B)$. \square

Theorem 3.6. In minimal structure, the arbitrary union of m - j open sets is always an m - j open set.

Proof. Let $A = \bigcup_i A_i$ be the arbitrary union of m - j open sets.

Hence $A \subseteq m_x\text{-Int}(m_x\text{-pcl}(A))$ which implies $\bigcup_i A_i \subseteq m_x\text{-Int}(m_x\text{-pcl}(\bigcup_i A_i))$. Therefore, each A_i is m - j open. Hence the arbitrary union of m - j open is always m - j open. \square

Remark 3.7. The intersection of any two m - j open sets need not be m - j open by the following example.

Consider $X = \{p, q, r\}$ with minimal structure m_x given by $m_x = \{\emptyset, X, \{p\}, \{s\}, \{q, r\}, \{p, q\}, \{p, r, s\}\}$. The corresponding m - j open set is given by $m\text{-}jO(X) = \{\emptyset, X, \{p\}, \{s\}, \{p, q\}, \{q, r\}, \{p, s\}, \{p, q, r\}, \{p, r, s\}, \{p, q, s\}, \{q, r, s\}\}$. Here $\{p, q\} \cap \{q, r\} = \{q\}$ which is not in the m - j open set.

Theorem 3.8. Every m - j open sets are m -preopen sets.

Proof. Let A be the m - j open set. Hence $A \subseteq m_x\text{-Int}(m_x\text{-pcl}(A))$. Since $\text{pcl}(A) \subseteq \text{Cl}(A)$ we have $A \subseteq m_x\text{-Int}(m_x\text{-Cl}(A))$. Hence A is m -preopen. \square

The converse of the above theorem is not true by the above example. $m\text{-}PO(X) = \{\emptyset, X, \{p\}, \{s\}, \{p, q\}, \{q, r\}, \{r, s\}, \{p, s\}, \{p, r\}, \{q, s\}, \{p, q, r\}, \{p, r, s\}, \{p, q, s\}, \{q, r, s\}\}$. Here $A = \{r, s\}$ is m -preopen but not m - j open.

Theorem 3.9. Every m - j open sets are m - b open sets.

Proof. We have to show that A is m - b open. Since A is m - j open, we have A is m -preopen. That is $A \subseteq m_x\text{-Int}(m_x\text{-Cl}(A))$. Hence we have $A \subseteq m_x\text{-Int}(m_x\text{-Cl}(A)) \cup m_x\text{-Cl}(m_x\text{-Int}(A))$. Therefore, A is m - b open. \square

The converse of the above theorem is not true by the following example. Consider $X = \{p, q, r\}$ with minimal structure m_x given by $m_x = \{\emptyset, X, \{p\}, \{q\}, \{p, r\}\}$ with $m\text{-}bO(X) = \{\emptyset, X, \{p\}, \{q\}, \{r\}, \{p, q\}, \{p, r\}\}$ and $m\text{-}jO(X) = \{\emptyset, X, \{p\}, \{q\}, \{p, q\}, \{p, r\}\}$. Here $A = \{r\}$ is m - b open but not m - j open.



4. m - j border, m - j frontier, m - j exterior of the sets

Definition 4.1. In a minimal structure (X, m_x) , $m_x-b_j(A) = A \setminus m_x-Int_j(A)$ is called the m - j border of A , where A is the subset of X .

Example 4.2. Consider the minimal structure (X, m_x) with $X = \{p, q, r, s\}$ and $m_x = \{\emptyset, X, \{p\}, \{s\}, \{q, r\}, \{p, q\}, \{p, r, s\}\}$. And the corresponding m - j open set is given by $m-jO(X) = \{\emptyset, X, \{p\}, \{s\}, \{p, q\}, \{q, r\}, \{p, s\}, \{p, q, r\}, \{p, r, s\}, \{p, q, s\}, \{q, r, s\}\}$. Let $A = \{r, s\}$ then $m_x-b_j(A) = \{r\}$.

Theorem 4.3. Consider the minimal structures (X, m_x) with the subset A of X , the following conditions are true:

- i) $m_x-b_j(A) \subseteq m_x-b(A)$
- ii) $m_x-Int_j(A) \cup m_x-b_j(A) = A$
- iii) $m_x-Int_j(A) \cap m_x-b_j(A) = \emptyset$
- iv) A is an m - j open set if and only if $m_x-b_j(A) = \emptyset$
- v) $m_x-b_j(m_x-Int_j(A)) = \emptyset$
- vi) $m_x-Int_j(m_x-b_j(A)) = \emptyset$
- vii) $m_x-b_j(m_x-b_j(A)) = m_x-b_j(A)$
- viii) $m_x-b_j(A) = A \cap m_x-Cl_j(X - A)$
- ix) $m_x-b_j(A) = m_x-D_j(X - A)$

Proof. i) $m_x-b_j(A) = A \setminus m_x-Int_j(A) \subseteq A \setminus m_x-Int(A) \subseteq m_x-b(A)$.
 ii) $m_x-Int_j(A) \cup m_x-b_j(A) = m_x-Int_j(A) \cup (A \setminus m_x-Int_j(A)) = A$.
 iii) $m_x-Int_j(A) \cap m_x-b_j(A) = m_x-Int_j(A) \cap (A \setminus m_x-Int_j(A)) = \emptyset$.
 iv) A is an m - j open set if and only if $m_x-Int_j(A) = A$ if and only if $A \setminus m_x-Int_j(A) = \emptyset$ if and only if $m_x-b_j(A) = \emptyset$.
 v) By the definition of m - j border, we have $m_x-b_j(A) = A \setminus m_x-Int_j(A)$. When $A = m_x-Int_j(A)$ we have $m_x-b_j(A) = m_x-Int_j(A) \setminus m_x-Int_j(A) = \emptyset$.
 vi) If $x \in m_x-Int_j(m_x-b_j(A))$ then $x \in m_x-b_j(A)$. On the other hand, since $m_x-b_j(A) \subset A$, we have $x \in m_x-Int_j(m_x-b_j(A)) \subset m_x-Int_j(A)$. Hence $x \in m_x-Int_j(A) \cap m_x-b_j(A)$ which is a contradiction to (iii). Thus $m_x-Int_j(m_x-b_j(A)) = \emptyset$.
 vii) Since $m_x-b_j(m_x-Int_j(A)) = \emptyset$, we have $m_x-b_j(m_x-b_j(A)) = m_x-b_j(A \setminus m_x-Int_j(A)) = m_x-b_j(A)$.
 viii) $m_x-b_j(A) = A \setminus m_x-Int_j(A) = A \setminus (X \setminus m_x-Cl_j(X - A)) = A \cap m_x-Cl_j(X - A)$.
 ix) $m_x-b_j(A) = A \setminus m_x-Int_j(A) = A \setminus (A \setminus m_x-D_j(X - A)) = m_x-D_j(A)$. \square

Consider the Example 4.2. If $A = \{p, s\}$ then $m_x-b_j(A) = \emptyset$ and $m_x-b(A) = \{s\}$. Hence $m_x-b(A) \not\subseteq m_x-b_j(A)$. In general, the converse part of the theorem 4.3(i) may not be true.

Definition 4.4. Let (X, m_x) be the minimal structure and A be the subset of X , then $m_x-Fr_j(A) = m_x-Cl_j(A) \setminus m_x-Int_j(A)$ is said to be m - j frontier of A .

Example 4.5. Let $X = \{a, b, c\}$ with minimal structure $m_x = \{\emptyset, X, \{a\}, \{b\}, \{b, c\}, \{a, b\}\}$ and the corresponding m - j open set is given by $m-jO(X) = \{\emptyset, X, \{a, b\}, \{b, c\}, \{a\}, \{b\}\}$. Let $A = \{a, b\}$ then $m_x-Fr_j(A) = \{c\}$.

Theorem 4.6. Consider the minimal structures (X, m_x) with the subset A of X the following conditions are true:

- i) $m_x-Fr_j(A) \subseteq m_x-Fr(A)$ where $Fr(A)$ denotes the frontier of A .
- ii) $m_x-Cl_j(A) = m_x-Int_j(A) \cup m_x-Fr_j(A)$.
- iii) $m_x-Int_j(A) \cap m_x-Fr_j(A) = \emptyset$.
- iv) $m_x-b_j(A) \subset m_x-Fr_j(A)$.
- v) $m_x-Fr_j(A) = m_x-b_j(A) \cup m_x-D_j(A)$.
- vi) A is an m - j open set if and only if $m_x-Fr_j(A) = m_x-D_j(A)$.
- vii) $m_x-Fr_j(A) = m_x-Cl_j(A) \cap m_x-Cl_j(X - A)$.
- viii) $m_x-Fr_j(A) = m_x-Fr_j(X - A)$.
- ix) $m_x-Fr_j(A)$ is m - j closed.
- x) $m_x-Fr_j(m_x-Fr_j(A)) \subseteq m_x-Fr_j(A)$.
- xi) $m_x-Fr_j(m_x-Int_j(A)) \subseteq m_x-Fr_j(A)$.
- xii) $m_x-Fr_j(m_x-Cl_j(A)) \subseteq m_x-Fr_j(A)$.
- xiii) $m_x-Int_j(A) = A \setminus m_x-Fr_j(A)$.

Proof. i) $m_x-Fr_j(A) = m_x-Cl_j(A) \setminus m_x-Int_j(A) \subseteq m_x-Cl(A) \setminus m_x-Int(A) \subseteq m_x-Fr(A)$
 ii) $m_x-Int_j(A) \cup m_x-Fr_j(A) = m_x-Int_j(A) \cup (m_x-Cl_j(A) \setminus m_x-Int_j(A)) = m_x-Cl_j(A)$
 iii) It is similar to (ii) and also by the definition.
 iv) $m_x-b_j(A) = A \setminus m_x-Int_j(A) \subseteq m_x-Cl_j(A) \setminus m_x-Int_j(A) \subseteq m_x-Fr_j(A)$.
 v) It follows from (ii) and from the fact that $m_x-Cl_j(A) = A \cup m_x-D_j(A)$.
 vi) It follows from (v) and the fact that if A is an m - j open set if and only if $m_x-b_j(A) = \emptyset$.
 vii) $m_x-Fr_j(A) = m_x-Cl_j(A) \setminus m_x-Int_j(A) = m_x-Cl_j(A) \cap m_x-Cl_j(X - A)$.
 viii) It is similar to (vii) and by the definition of m - j frontier of A .
 ix) $m_x-Cl_j(m_x-Fr_j(A)) = m_x-Cl_j(m_x-Cl_j(A) \cap m_x-Cl_j(X - A)) = m_x-Cl_j(m_x-Cl_j(A)) \cap m_x-Cl_j(m_x-Cl_j(X - A)) = m_x-Fr_j(A)$.
 x) $m_x-Fr_j(m_x-Fr_j(A)) = m_x-Cl_j(m_x-Fr_j(A)) \cap m_x-Cl_j(X \setminus m_x-Fr_j(A)) \subseteq m_x-Cl_j(m_x-Fr_j(A)) = m_x-Fr_j(A)$.
 xi) It is similar to (x).
 xii) $m_x-Fr_j(m_x-Cl_j(A)) = m_x-Cl_j(m_x-Cl_j(A)) \setminus m_x-Int_j(m_x-Cl_j(A)) = m_x-Cl_j(A) \setminus m_x-Int_j(m_x-Cl_j(A)) = m_x-Cl_j(A) \setminus m_x-Int_j(A) = m_x-Fr_j(A)$.
 xiii) $A \setminus m_x-Fr_j(A) = A \setminus (m_x-Cl_j(A) \setminus m_x-Int_j(A)) = m_x-Int_j(A)$. \square

The converse part of (i) and (iv) of the above theorem are not true by the following example. Consider $X = \{p, q, r, s\}$ with minimal structure given by $m_x = \{\emptyset, X, \{p\}, \{s\}, \{q, r\}, \{p, q\}, \{p, r, s\}\}$. The corresponding m - j open set is given by $m-jO(X) = \{\emptyset, X, \{p\}, \{s\}, \{p, q\}, \{q, r\}, \{p, s\}, \{p, q, r\}, \{p, r, s\}, \{p, q, s\}, \{q, r, s\}\}$. If $A = \{p, r\}$ then $m_x-Fr(A) = \{b, c\} \not\subseteq \{c\} = m_x-Fr_j(A)$. Consider $X = \{p, q, r\}$ with minimal structure given by $m_x = \{\emptyset, X, \{p\}, \{q\}, \{q, r\}, \{p, q\}\}$ and the corresponding m - j open set is given by $m-jO(X) = \{\emptyset, X, \{p, q\}, \{q, r\}, \{p\}, \{q\}\}$. If $A = \{a, b\}$, then $m_x-Fr_j(A) = \{c\}$ and $m_x-b_j(A) = \emptyset$. Hence $m_x-Fr_j(A) \not\subseteq m_x-b_j(A)$.



Definition 4.7. Let A be the subset of minimal structure (X, m_x) , then $m_x\text{-Ext}_j(A) = m_x\text{-Int}_j(X \setminus A)$ is said to be m - j exterior of A .

Example 4.8. Consider $X = \{p, q, r\}$ and $m_x = \{\emptyset, X, \{p\}, \{r\}, \{p, r\}\}$ and the corresponding m - j open set is given by $m_jO(X) = \{\emptyset, X, \{p\}, \{q\}, \{r\}, \{p, q\}, \{p, r\}\}$. Let $A = \{q\}$, then $m_x\text{-Ext}_j(A) = \{p, r\}$.

Theorem 4.9. For the subset A and B of X , the following conditions are true

- i) $m_x\text{-Ext}(A) \subseteq m_x\text{-Ext}_j(A)$ where $\text{Ext}(A)$ denotes the exterior of A .
- ii) $m_x\text{-Ext}_j(A)$ is open.
- iii) $m_x\text{-Ext}_j(A) = m_x\text{-Int}_j(X \setminus A) = X \setminus m_x\text{-Cl}_j(A)$.
- iv) $m_x\text{-Ext}_j(m_x\text{-Ext}_j(A)) = m_x\text{-Int}_j(m_x\text{-Cl}_j(A))$.
- v) If $A \subseteq B$, then $m_x\text{-Ext}_j(A) \subseteq m_x\text{-Ext}_j(B)$.
- vi) $m_x\text{-Ext}_j(A \cup B) \subseteq m_x\text{-Ext}_j(A) \cap m_x\text{-Ext}_j(B)$.
- vii) $m_x\text{-Ext}_j(A) \cup m_x\text{-Ext}_j(B) \subseteq m_x\text{-Ext}_j(A \cap B)$.
- viii) $m_x\text{-Ext}_j(X) = \emptyset$.
- ix) $m_x\text{-Ext}_j(\emptyset) = X$.
- x) $m_x\text{-Ext}_j(A) = m_x\text{-Ext}_j(X \setminus m_x\text{-Ext}_j(A))$.
- xi) $m_x\text{-Int}_j(A) = m_x\text{-Ext}_j(m_x\text{-Ext}_j(A))$.
- xii) $X = m_x\text{-Int}_j(A) \cup m_x\text{-Ext}_j(A) \cup m_x\text{-Fr}_j(A)$.

Proof. i) Since every interior point of A is the j -interior of A in minimal structure, we have $m_x\text{-Int}(A) \subseteq m_x\text{-Int}_j(A)$. Therefore, $m_x\text{-Ext}(A) = m_x\text{-Int}(X \setminus A) \subseteq m_x\text{-Int}_j(X \setminus A) = m_x\text{-Ext}_j(A)$.

ii) Since $m_x\text{-Ext}_j(A) = m_x\text{-Int}_j(X \setminus A)$, then $m_x\text{-Ext}_j(A)$ is open.

iii) It is obvious that $m_x\text{-Ext}_j(A) = m_x\text{-Int}_j(X \setminus A) = X \setminus m_x\text{-Cl}_j(A)$.

iv) Since $m_x\text{-Int}_j(X \setminus A) = X \setminus m_x\text{-Cl}_j(A)$, we have $m_x\text{-Ext}_j(m_x\text{-Ext}_j(A)) = m_x\text{-Ext}_j(m_x\text{-Int}_j(X \setminus A)) = m_x\text{-Int}_j(X \setminus m_x\text{-Int}_j(X \setminus A)) = m_x\text{-Int}_j(m_x\text{-Cl}_j(A))$.

v) Since $A \subseteq B$, we have $m_x\text{-Int}_j(A) \subseteq m_x\text{-Int}_j(B)$. Therefore by definition we have $m_x\text{-Ext}_j(A) \subseteq m_x\text{-Ext}_j(B)$.

vi) Using $m_x\text{-Int}_j(A) \cup m_x\text{-Int}_j(B) \subseteq m_x\text{-Int}_j(A \cup B)$, we have $m_x\text{-Ext}_j(A \cup B) = m_x\text{-Int}_j(X - (A \cup B)) = m_x\text{-Int}_j((X - A) \cap (X - B)) \subseteq m_x\text{-Int}_j(X - A) \cap m_x\text{-Int}_j(X - B) = m_x\text{-Ext}_j(A) \cap m_x\text{-Ext}_j(B)$.

vii) The proof is similar to (vi).

viii) $m_x\text{-Ext}_j(X) = m_x\text{-Int}_j(X^c) = \emptyset$.

ix) $m_x\text{-Ext}_j(\emptyset) = m_x\text{-Int}_j(\emptyset^c) = X$.

x) Since $m_x\text{-Int}_j(m_x\text{-Int}_j(A)) = m_x\text{-Int}_j(A)$, by the definition of j -exterior of A we have $m_x\text{-Ext}_j(A) = m_x\text{-Ext}_j(X \setminus m_x\text{-Ext}_j(A))$.

xi) Since $m_x\text{-Ext}_j(A) = m_x\text{-Int}_j(X \setminus A)$, we have $m_x\text{-Int}_j(A) = m_x\text{-Ext}_j(m_x\text{-Ext}_j(A))$.

xii) It is obvious from the definition that $X = m_x\text{-Int}_j(A) \cup m_x\text{-Ext}_j(A) \cup m_x\text{-Fr}_j(A)$. \square

- [2] I. Arockiarani and D. Sasikala, A new class of j -separations is generalized Topological spaces, *Asian Journal of Current Engineering and Maths* 1-4(2012), 188–191.
- [3] M. Caldas, M. Gangster, F. Jafari, P. Noiri and N. Rajesh, Properties of α open sets in ideal minimal spaces, *Sci. Stud. Res. Ser. Math. Inform.*, 27(2)(2017), 83–100.
- [4] Ennis Rosas and Neelamegarajan Rajesh, Some new types of open and closed sets in minimal structures-II, *International Mathematical Forum*, 44(2009), 2185–2198.
- [5] Jafari, N. Rajesh and R. Saranya, Semi-open sets in ideal minimal spaces, *Vasile Alecsandri University of Bacau Faculty of Sciences Scientific Studies and Research Series Mathematics and Informatics*, 27(1)(2017), 33-48.
- [6] Miguel Calidas, A note on some application of α open sets, 2003, 125-130.
- [7] W. K. Min, m -semiopen sets and m -semicontinuous functions on spaces with minimal structure, *Honam Mathi.*, (31)(2)(2009), 239–245.
- [8] W. K. Min and Y.K.Kim, m -preopen sets and m -precontinuity on spaces with minimal structures, *Advances in Fuzzy Sets and Systems*, (4)(3)(2009), 27–245.
- [9] W. K. Min, αm -open sets and αm -continuous functions, *Commun. Korean Math. Soc.*, (25)(2)(2010), 251–256.
- [10] Mohammed S. Sarsak, On b -open sets and associated generalized open sets, *Questions and Answers in General Topology*, 27(2009), 157–173.
- [11] Norman Levine, Semi-open sets and semi-continuity in topological spaces, *The American Mathematical Monthly*, 70(1)(1963), 36–41.
- [12] V. Popa and T. Noiri, On m -continuous functions and product spaces, *Annales Univ. Sci.*, 47(2004), 65-89.
- [13] Sucharita Chakrabarti, On m -baire spaces, *International Mathematical Forum*, 6(2)(2011), 95–101.
- [14] Young Bae Jun, Scong Woo Jeong, Hyeon Jeong Lee and Joon Woon Lee, Applications of pre-open sets, *Universitat Politècnica de València*, 9(2)(2008), 213–228.

ISSN(P):2319 – 3786

Malaya Journal of Matematik

ISSN(O):2321 – 5666

References

- [1] M.H.H. Ameer, Pre-open sets in minimal bitopological spaces, *Journal of Kufa for Mathematics and Computer*, 2(3)(2015), 26–42.

